



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Fixed Partial Prosthesis

Classification and clinical rationale

Acknowledgements to Dr Matsubara

Learning outcomes



- Understand and discuss the advantages and disadvantages of different types of FPD
- Discuss the differences between the types of bridges
- Discuss tooth preparation for bonded bridges
- Discuss fibre-reinforced bonded bridges
- Discuss the concept of Shortened dental arch
- Discuss the clinical rationale of prosthetic treatment using FPD: factors that affect the prognosis of the treatment

Practical advice for successful clinical treatment with
resin-bonded bridges

Reading:

Stephanie King,¹ Banoo Sood¹ and Martin P. Ashley^{*2}

Classification


PROSTHODONTICS



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Material

- Acrylic, composite (temporary bridge)
- Cast metal: gold alloy
- Metal-ceramic (PFM)
- Resin-Veneered
- All ceramic (e.g. zirconia)



Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia


Cement-Retained FPD

Metal-ceramic



- Cemented with conventional luting material (GIC, RMGIC, Zinc phosphate)

All ceramic



- Cemented with conventional luting material
- Bonded with resin cement

Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

Early bonded FPD



Extracted natural teeth
Acrylic teeth
(temporary restoration)



70s - Rochette bridge

Etched cast



Maryland Bridge

Classification



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia

Resin-bonded FPD

Traditional approach

ADVANTAGES

- Minimal removal of tooth structure (wing thickness 0.8mm)
- Supragingival preparation, easy impression making
- Interim restoration not usually required

Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

Traditional approach



ADVANTAGES

- Minimal removal of tooth structure
- Supragingival preparation, easy impression making
- Interim restoration not usually required

DISADVANTAGES

- Reduced restoration longevity
- Space correction is limited
- Good alignment of abutment teeth is required
- Esthetics is compromised on anterior

Classification



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia

Resin-bonded FPD

Traditional approach

INDICATIONS

- Short edentulous space (one tooth)
- Healthy/Unrestored abutments (enamel surface)
- Significant clinical crown length
- Temporary restoration - space maintenance
- Open bite

Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

Traditional approach

INDICATIONS

- Short edentulous space (one tooth)
- Healthy/Unrestored abutments (enamel surface)
- Significant clinical crown length
- Temporary restoration - space maintenance
- Open bite

CONTRAINDICATIONS

- Damaged abutments
- Large edentulous area
- Unfavorable angulation and rotation of abutments
- Parafunctional habits or heavy occlusal forces
- Deep bite
- Mobile teeth
- Alloy allergy

Classification



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

Traditional approach

Design and construction:


- Cantilevered design (anterior)

Research | Published: 10 April 2015

Survival characteristics of 771 resin-retained bridges provided at a UK dental teaching hospital

[P. A. King](#), [L. V. Foster](#), [R. J. Yates](#), [R. G. Newcombe](#) & [M. J. Garrett](#)

Figure 3



survival curves for 476 cantilever bridges and a group comprising 292 fixed-fixed and
jes

Resin-bonded FPD


Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
of Western Australia



Stability
&
Retention

Single path of Insertion



Resin-bonded FPD

Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
of Western Australia



Guide planes: maximum surface area of enamel contact, POI, Resistance against horizontal movement

Axial grooves
Resistance against horizontal movement



Rest seat
Resistance against vertical occlusal forces

Resin-bonded FPD



Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Resin-bonded FPD


Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Resin-bonded FPD




Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Resin-bonded FPD





Metal-ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
of Western Australia



Resin-bonded FPD

Metal-ceramic



Resin-bonded FPD

All ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Resin-bonded FPD

All ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Resin-bonded FPD




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

PRINCIPLES OF TOOTH PREPARATION

- Intra-enamel preparation
- Supragingival margins
- Maximize surface area for retainers
- Bracing of abutments
- Single path of insertion
- Clearance for retainers
- Rest seats



Resin-bonded FPD

All ceramic



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Retainers:

Inlay

Onlay



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

All ceramic



Pegoraro et al 2013



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Resin-bonded FPD

All ceramic



Pegoraro et al 2013

Resin-bonded FPD

Fiber-reinforced




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia


Fiber-reinforced bonded bridge: natural or acrylic tooth



Ribbond®




Leno weave



Woven



Braided



Unidirectional

Types of fibers:

- Glass
- Polyethylene
- Kevlar

Resin-bonded FPD

Fiber-reinforced




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Fiber-reinforced bonded bridge



Resin-bonded FPD

Fiber-reinforced



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



THE UNIVERSITY OF
WESTERN
AUSTRALIA

Fiber-reinforced bonded bridge



Resin-bonded FPD

Fiber-reinforced



THE UNIVERSITY OF
WESTERN
AUSTRALIA





Oral Health Centre
of Western Australia



THE UNIVERSITY OF
WESTERN
AUSTRALIA

Fiber-reinforced bonded bridge



Resin-bonded FPD

Fiber-reinforced



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Fiber-reinforced bonded bridge



Resin-bonded FPD

Fiber-reinforced



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**

Fiber-reinforced bonded bridge




Implant Supported Bridge



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



Implant Supported Bridge



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



CLINICAL RATIONALE



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia



CLINICAL RATIONALE



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Consequences of tooth loss

- Loss of masticatory efficiency – affect systemic health
- Loss of quality, width and height of bone
- Loss of soft tissue volume
- Aesthetics is compromised
- Phonetic is compromised (dental consonants)
- Occlusal instability

CLINICAL RATIONALE





THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Consequences of tooth loss

- Occlusal changes: Interferences, unbalanced muscle contraction, temporomandibular disorders (TMD)



CLINICAL RATIONALE



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
of Western Australia

Consequences of tooth loss

- Occlusal changes: Interferences, unbalanced muscle contraction, temporomandibular disorders (TMD)

Shortened dental arch concept



Witter et al. 1999

CLINICAL RATIONALE



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Rationale for Intervention:

- Is there a need for improved aesthetics?
- Does the patient need to improve masticatory function ?
- Is there a need for improved occlusal stability?

Need vs Demand



CLINICAL RATIONALE



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
Perth, Australia

Rationale for Intervention:

Aesthetics:

- What the patient wants
- What the patient needs
- What the patient can get

What is the patient's expectation?
Can it be achieved?
What is technically possible?



CLINICAL RATIONALE




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Aesthetics
Lip support, high smile line



CLINICAL RATIONALE



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Clinical Rationale of Tooth Supported Bridgework



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

NEVER OVERLOAD ABUTMENT TEETH!

- Direction of Occlusal forces:
 - 1) Mandibular stability
 - 2) Axial occlusal load
 - 3) During lateral excursions: no interference in the working side
 - 4) During lateral excursions: disocclusion in the non-working side
 - 5) During protrusion: disocclusion of posterior teeth
- Intensity of Occlusal Forces:
 - Parafunctional habits
 - Age, Gender

Clinical Rationale of Tooth Supported Bridgework



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

Assessing potential abutment teeth:

- Crown/Root Ratio
- Root configuration
- Periodontal Ligament Area

Ante's Law: "Root surface area of the abutment teeth has to equal or surpass that of the teeth being replaced with pontics"





Fig 7-12 The combined root surface area of the canine and second molar ($A_c + A_{2m}$) is exceeded by that of the teeth being replaced ($A_{1p} + A_{2p} + A_{1m}$). A fixed partial denture would be a poor choice in this situation.




Fig 7-10 The combined root surface area of the second premolar and the second molar ($A_{2p} + A_{2m}$) is greater than that of the first molar being replaced (A_{1m}).

Clinical Rationale of Tooth Supported Bridgework




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Crown-to-root ratio

Minimum 1:1



Shillingburg 1997

Clinical Rationale of Tooth Supported Bridgework




THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Root shape and angulation



Clinical Rationale of Tooth Supported Bridgework





THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia

- Root surface area



JEPSEN A. Root surface measurement and a method for x-ray determination of root surface area. Acta Odontol Scand. 1963.

Clinical Rationale of Tooth Supported Bridgework



THE UNIVERSITY OF
WESTERN
AUSTRALIA




Oral Health Centre
of Western Australia

- Length of edentulous area


Ante's Law

The total root surface area (periodontal membrane) of the abutment teeth for a FPD must be equal to or greater than the root surface area of the teeth being replaced


(Irvin Ante, 1926)



1



2



3

Clinical Rationale of Tooth Supported Bridgework





THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Length of edentulous area



- Fracture of a porcelain veneer
- Breakage of a connector
- Loosening of a retainer
- Unfavorable soft tissue response
- Fracture of abutment

Clinical Rationale of Tooth Supported Bridgework





THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Dental arch shape



- Opposing arch

Clinical Rationale of Tooth Supported Bridgework



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Dental arch shape




Fig 7-20 Secondary retention (R) must extend a distance from the primary interabutment axis equal to the distance that the pontic lever arm (P) extends in the opposite direction.

Fundamentals of Fixed Prosthodontics




Fig 7-19 The retainers on secondary abutments will be placed in tension when the pontics flex, with the primary abutments acting as fulcrums.

Clinical Rationale of Tooth Supported Bridgework



THE UNIVERSITY OF
WESTERN
AUSTRALIA



Oral Health Centre
of Western Australia

- Direction of Occlusal forces
- Intensity of Occlusal Forces
- Assess root surface area
- Length of edentulous area
- Root shape and angulation
- Crown-to-root ratio
- Dental arch shape
- Opposing arch

Thanks for listening!



THE UNIVERSITY OF
**WESTERN
AUSTRALIA**



Oral Health Centre
of Western Australia
